



Recombinant Protein Technical Manual

Recombinant Human CNDP1 Protein (His Tag)(Active)
RPES2304

Product Data:

Product SKU: RPES2304

Size: 10 μ g

Species: Human

Expression host: HEK293 Cells

Uniprot: NP_116038.4

Protein Information:

Molecular Mass: 55.3 kDa

AP Molecular Mass: 60-65 kDa

Tag: C-His

Bio-activity: Measured by its ability to cleave carnosine (β -Ala-L-His) in a two-step assay. The specific activity is > 250 pmoles/min/ μ g.

Purity: > 90 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per μ g of the protein as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Beta-Ala-His Dipeptidase; CNDP Dipeptidase 1; Carnosine Dipeptidase 1; Glutamate Carboxypeptidase-Like Protein 2; Serum Carnosinase; CNDP1; CN1; CPGL2;HsT2308

Immunogen Information:

Sequence: Ser27-His507

Background:

CNDP1, also known as carnosine dipeptidase 1, glutamate carboxypeptidase-like protein 2 (CPGL-2) or carnosinase 1 (CN1), is a member of the M20 metalloprotease family. The CNDP1 gene contains trinucleotide (CTG) repeat length polymorphism in the coding region, which has been demonstrated to be associated with susceptibility to developing diabetic nephropathy, for carnosine protection against the adverse effects of high glucose levels on renal cells. In humans, CNDP1 is secreted from the liver into the serum. In other mammals, including rodents, CNDP1 is expressed exclusively within the kidney and lacks a signal peptide. CNDP1 protein is a secreted homodimeric dipeptidase that specifically hydrolyzes L-carnosine (β -alanyl-L-histidine), and is identified as human carnosinase expressed in the brain. CNDP1 has been associated with diabetic nephropathy in Europeans and European Americans, but not African-Americans. It was identified and confirmed as a risk factor, were cross-sectional and mostly in patients with type 2 diabetes. The polymorphisms of CNDP1 can be excluded as a risk factor for nephropathy in type 1 diabetes. In addition, CNDP1 is also suggested to be implicated in the actions of neuroprotection and neurotransmiting.