



Recombinant Protein Technical Manual
Recombinant Cynomolgus B7/CD80 Protein (Fc Tag)
RPES2654

Product Data:

Product SKU: RPES2654

Size: 10µg

Species: Cynomolgus

Expression host: Human Cells

Uniprot: G7NXN7

Protein Information:

Molecular Mass: 51 kDa

AP Molecular Mass: 70-90 kDa

Tag: C-Fc

Bio-activity:

Purity: > 95% as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of 50 mM Tris, 100 mM Glycine, pH 7.5.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: T-lymphocyte activation antigen CD80; Activation B7 antigen; B7; CD80

Immunogen Information:

Sequence: Val35-Asn242

Background:

Cynomologous Cluster of Differentiation 80, also called B7, is a member of cell surface immunoglobulin superfamily. It is expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. CD80 plays key, yet distinct roles in the activation of T cells. B7/CD80 and B7-2/CD86, together with their receptors CD28 and CTLA4, constitute one of the dominant co-stimulatory pathways that regulate T- and B- cell responses. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells, macrophages and dendritic cells. Although both CTLA-4 and CD28 can bind to the same ligands, CTLA-4 binds to B7 and B7-2 with a 2000 fold higher affinity than CD28 and is involved in the down-regulation of the immune response. CD80 is thus regarded as promising therapeutic targets for autoimmune diseases and various carcinomas.