

Recombinant Protein Technical Manual Recombinant Cynomolgus TIM-3/HAVCR2 Protein (Fc Tag) RPES2738

## **Product Data:**

Product SKU: RPES2738

Species: Cynomolgus

Size:  $10 \mu g$ 

Expression host: Human Cells

Uniprot: G7P6Q7

## **Protein Information:**

Molecular Mass:	46.3 kDa
AP Molecular Mass:	55-75 kDa
Tag:	C-Fc
Bio-activity:	
Purity:	> 95% as determined by reducing SDS-PAGE.
Endotoxin:	< 1.0 EU per $\mu g$ as determined by the LAL method.
Storage:	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH7.4.
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	T cell immunoglobulin and mucin domain3; HAVCR2; Tim-3; TIM3

## Sequence: Ser22-Arg201

## Background:

T cell immunoglobulin and mucin domain 3 is a member of the TIM family of immune regulating molecules. Mature cynomolgus TIM3 consists of a 182 amino acid (aa)extracellular domain (ECD), a 21 aa transmembrane segment, and a 78 aa cytoplasmic tail. TIM3 is up-regulated on several populations of activated myeloid cells (macrophage, monocyte, dendritic cell, microglia, mast cell) and T cells (Th1, CD8+, NK, Treg). Its binding to Galectin9 induces a range of immunosuppressive functions which enhance immune tolerance and inhibit anti-tumor immunity. TIM3 ligation attenuates CD8+ and Th1 cell responses and promotes the activity of Treg and myeloid derived suppressor cells. TIM3 interactions with Galectin-9 can trigger immune stimulatory effects, such as the coactivation of NK cell cytotoxicity.