



Recombinant Protein Technical Manual
Recombinant Mouse GLO1/Glyoxalase 1 Protein (His
Tag)
RPES2914

Product Data:

Product SKU: RPES2914

Size: 20µg

Species: Mouse

Expression host: E. coli

Uniprot: NP_079650.3

Protein Information:

Molecular Mass: 21.6 kDa

AP Molecular Mass: 25 & 48 kDa

Tag: N-His

Bio-activity:

Purity: > 95 % as determined by SDS-PAGE

Endotoxin: Please contact us for more information.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: 0610009E22Rik;1110008E19Rik;2510049H23Rik;AW550643;Glo;Glor;Glos;Glo1-r;Glo1-s;GLY1;Qglo

Immunogen Information:

Sequence: Ala 2-Ile 184

Background:

Lactoylglutathione lyase, also known as Methylglyoxalase, Aldoketomutase, Glyoxalase I, Ketone-aldehyde mutase, S-D-lactoylglutathione methylglyoxal lyase and GLO1, is a member of the glyoxalase I family. GLO1 / Glyoxalase I is a ubiquitous cellular defense enzyme involved in the detoxification of methylglyoxal, a cytotoxic byproduct of glycolysis. Accumulative evidence suggests an important role of GLO1 expression in protection against methylglyoxal-dependent protein adduction and cellular damage associated with diabetes, cancer, and chronological aging. GLO1 / Glyoxalase I has been implicated in anxiety-like behavior in mice and in multiple psychiatric diseases in humans. GLO1 / Glyoxalase I catalyzes the conversion of hemimercaptal, formed from methylglyoxal and glutathione, to S-lactoylglutathione. GLO1 / Glyoxalase I exists in three separable isoforms which originate from two alleles in the genome. These correspond to two homodimers and one heterodimer composed of two subunits showing different electrophoretic properties. GLO1 upregulation may play a functional role in glycolytic adaptations of cancer cells.