

Recombinant Protein Technical Manual Recombinant Mouse SR-BI/CD36L1 Protein (His Tag) RPES2926

Product Data:

Product SKU: RPES2926

Species: Mouse

Size: 20µg

Expression host: HEK293 Cells

Uniprot: NP_058021.1

Protein Information	

Molecular Mass:	47 kDa
AP Molecular Mass:	70-90 kDa
Tag:	C-His
Bio-activity:	
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	AI120173;CD36;Cd36l1;Cla;Cla1;D5Ertd460e;Hdlq1;Hlb398;mSR-BI;SR-B1;SR- BI;Srb1;SRBI

Sequence: Pro 33-Val 440

Background:

Scavenger receptor class B, member 1 (SCARB1), also known as CD36L1, is a member of the scavenger receptor family. SCARB1 is expressed primarily in liver and non placental steroidogenic tissues, and predominantly localized to cholesterol and sphingomyelin-enriched domains within the plasma membrane. SCARB1 is proposed as a receptor for different ligands such as phospholipids, cholesterol ester, lipoproteins, phosphatidylserine and apoptotic cells, and is involved in a wide variety of physilogical processes. As a key component in the reverse cholesterol transport pathway, SCARB1 binds high density lipoproteins (HDLs) and mediates selective cholesterol uptake by a mechanism distinct from the LDL pathway. High density lipoproteins (HDLs) play a critical role in cholesterol metabolism and their plasma concentrations are inversely correlated with risk for atherosclerosis. SCARB1 may thus serve as a useful marker that predicts variation in baseline lipid levels and postprandial lipid response. The mouse SCARB1 has been shown to exert actions in determining the levels of plasma lipoprotein cholesterol and the accumulation of cholesterol stores in the adrenal gland.