



# Recombinant Protein Technical Manual

## Recombinant Human FGFR3/CD333 Protein (alpha IIIb, His Tag)

RPES2977

### Product Data:

**Product SKU:** RPES2977

**Size:** 50µg

**Species:** Human

**Expression host:** HEK293 Cells

**Uniprot:** NP\_001156685.1

### Protein Information:

**Molecular Mass:** 40 kDa

**AP Molecular Mass:**

**Tag:** C-His

**Bio-activity:**

**Purity:** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from sterile PBS, pH 7.4

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:**

**Synonyms:** Fibroblast growth factor receptor 3; FGFR-3;FGFR3; JTK4; IIIc;ACH;CD333;CEK2;HSFGFR3EX

## Immunogen Information:

**Sequence:** Met 1-Gly377

## Background:

FGFR3, also known as CD333, is a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid sequence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. FGFRs are transmembrane catalytic receptors that have intracellular tyrosine kinase activity. Mutations in FGFR genes are the cause of several human developmental disorders characterized by skeletal abnormalities such as achondroplasia, and upregulation of FGFR expression may lead to cell transformation and cancer. FGFR3, a full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR3 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. FGFR3 binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in FGFR3 gene lead to craniosynostosis and multiple types of skeletal dysplasia. Three alternatively spliced transcript variants that encode different protein isoforms have been described. CD333 is the receptor for acidic and basic fibroblast growth factors.