



Recombinant Protein Technical Manual

Recombinant Cynomolgus B2M/Beta-2-microglobulin Protein (His Tag)
RPES3053

Product Data:

Product SKU: RPES3053

Size: 10µg

Species: Cynomolgus

Expression host: Human Cells

Uniprot: Q8SPW0

Protein Information:

Molecular Mass: 12.5 kDa

AP Molecular Mass: 14 kDa

Tag: C-His

Bio-activity:

Purity: > 95% as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Beta-2-Microglobulin; B2M; β2-Microglobulin

Immunogen Information:

Sequence: Ile21-Met119

Background:

β -2-Microglobulin (B2M) is a secreted protein with 1 Ig-like C1-type (immunoglobulin-like) domain which belongs to the beta-2-microglobulin family. B2M component of major histocompatibility complex (MHC) class I molecules, involved in the presentation of peptide antigens to the immune system. Polymers of beta 2-microglobulin can be found in tissues from patients on long-term hemodialysis. B2M is a protein found on the surface of many cells and plentiful on the surface of white blood cells. Serum B2M concentration is increased in renal diseases, various malignant diseases and some inflammatory and autoimmune disorders. B2M may adopt the fibrillar configuration of amyloid in certain pathologic states. The capacity to assemble into amyloid fibrils is concentration dependent. B2M has been shown as a marker for monitoring inflammatory disease activity and it appears likely to have a destructive role in amyloidosis-related arthritis. B2M might be involved in the OA (osteoarthritis) pathogenesis. Defects in B2M are the cause of hypercatabolic hypoproteinemia. Affected individuals show marked reduction in serum concentrations of immunoglobulin and albumin, probably due to rapid degradation. B2M could be a potential therapeutic target in ovarian cancer.