



Recombinant Protein Technical Manual

**Recombinant Mouse Pleiotrophin/PTN/HB-GAM
Protein (Fc Tag)**
RPES3278

Product Data:

Product SKU: RPES3278

Size: 50µg

Species: Mouse

Expression host: HEK293 Cells

Uniprot: P63089

Protein Information:

Molecular Mass: 43.7 kDa

AP Molecular Mass: 47 kDa

Tag: N-Fc

Bio-activity:

Purity: > 85 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Pleiotrophin; PTN; Heparin-binding brain mitogen; HBBM; Heparin-binding growth factor 8; HBGF-8; Osteoblast-specific factor 1; OSF;HARP;HB-GAM;HBBN;HBNF;OSF;Osf;Osf1

Immunogen Information:

Sequence: Gly 33-Asp 168

Background:

HB-GAM belongs to the pleiotrophin family. During embryonic and early postnatal development, HB-GAM is expressed in the central and peripheral nervous system and also in several non-neural tissues, notably lung, kidney, gut and bone. While in the adult central nervous system, it is expressed in an activity-dependent manner in the hippocampus where it can suppress long term potentiation induction. HB-GAM has a low expression in other areas of the adult brain, but it can be induced by ischemic insults, or targeted neuronal damaged in the entorhinal cortex or in the substantia nigra pars compacta. It is structurally related to midkine and retinoic acid induced heparin-binding protein and has a high affinity for heparin. HB-GAM binds anaplastic lymphoma kinase (ALK) which induces MAPK pathway activation, an important step in the anti-apoptotic signaling of PTN and regulation of cell proliferation. It also functions as a secreted growth factor and induces neurite outgrowth and which is mitogenic for fibroblasts, epithelial, and endothelial cells.