



# Recombinant Protein Technical Manual

**Recombinant Human CD226/DNAM Protein (His Tag)(Active)**  
RPES3693

## Product Data:

**Product SKU:** RPES3693

**Size:** 50µg

**Species:** Human

**Expression host:** HEK293 Cells

**Uniprot:** NP\_006557.2

## Protein Information:

**Molecular Mass:** 27.6 kDa

**AP Molecular Mass:** 40-50 kDa

**Tag:** C-His

**Bio-activity:** Measured by its binding ability in a functional ELISA. Immobilized human CD226 at 2 µg/ml (100 µl/well) can bind human CD112 with a linear rangel of 6.4-800 ng/ml.

**Purity:** > 97 % as determined by reducing SDS-PAGE.

**Endotoxin:** < 1.0 EU per µg as determined by the LAL method.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from sterile PBS, pH 7.4

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:** Functional ELISA

**Synonyms:** DNAM;DNAM1;PTA1;TLISA1

## Immunogen Information:

**Sequence:** Met 1-Asn 247

## Background:

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD226, also known as PTA1 or DNAM, is a member of the immunoglobulin superfamily containing 2 Ig-like domains of the V-set. High rate of CD226 (Cluster of Differentiation 226) is found on the surface of natural killer cells, platelets, monocytes and a subset of T cells. CD226 have binding sites with CD112 and CD155 and mediate cellular adhesion to other cells containing its ligands.