

Recombinant Protein Technical Manual Recombinant Human FGF/FGFa Protein

RPES3780

Product Data:

Product SKU: RPES3780 **Size:** 10μg

Species: Human Expression host: E. coli

Uniprot: P05230

Protein Information:

Molecular Mass: 17.3 kDa

AP Molecular Mass: 17 kDa

Tag:

Bio-activity:

Purity: > 95 % as determined by reducing SDS-PAGE.

Endotoxin: $< 1.0 \text{ EU per } \mu\text{g}$ as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Fibroblast Growth Factor 1; FGF; Acidic Fibroblast Growth Factor; aFGF;

Endothelial Cell Growth Factor; ECGFHeparin-Binding Growth Factor 1; HBGF;

FGF1; FGFA

Immunogen Information:

Sequence: Ala2-Asp155

Background:

FGF acidic, also known as ECGF, FGFand HBGF, is a non-glycosylated heparin binding growth factor that is expressed in the brain, kidney, retina, smooth muscle cells, bone matrix, osteoblasts, astrocytes and endothelial cells. It is a mitogenic peptide that is produced by multiple cell types and stimulates the proliferation of cells of mesodermal, ectodermal, and endodermal origin. Its association with heparan sulfate is a prerequisite for activation of FGF receptors. Internalized FGF acidic migrates to the nucleus where it is phosphorylated by nuclear PKC delta, exported to the cytosol, dephosphorylated, and degraded. Intracellular FGF acidic inhibits p53 activity and proapoptotic signaling.