

# Recombinant Protein Technical Manual Recombinant Rhesus macaque CD47/IAP Protein (Fc Tag) RPES3834

### **Product Data:**

**Product SKU:** RPES3834 **Size:** 10μg

Species: Rhesus macaque Expression host: Human Cells

Uniprot: F7A802

### **Protein Information:**

Molecular Mass: 40.8 kDa

AP Molecular Mass: 60-90 kDa

Tag: C-Fc

**Bio-activity:** 

**Purity:** > 95% as determined by reducing SDS-PAGE.

**Endotoxin:**  $< 1.0 \text{ EU per } \mu\text{g}$  as determined by the LAL method.

**Storage:** Lyophilized protein should be stored at < -20°C, though stable at room

temperature for 3 weeks. Reconstituted protein solution can be stored at  $4-7^{\circ}$ C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from a 0.2 μm filtered solution of 50 mM Tris, 100 mM Glycine, pH 7.5.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:** 

**Synonyms:** Leukocyte Surface Antigen CD47; Antigenic Surface Determinant Protein OA3;

Integrin-Associated Protein; IAP; Protein MER6; CD47; MER6

# Immunogen Information:

Sequence: Gln19-Pro139

## **Background:**

CD47(Integrin-Associated Protein,IAP) is a 40 ? 60 kDa variably glycosylated atypical member of the immunoglobulin superfamily. The ubiquitously expressed CD47 binds to SIRP family members on macrophages, neutrophils, and T cells. CD47 is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The protein is also a receptor for the C-terminal cell-binding domain of thrombospondin, and it may play a role in membrane transport and signal transduction. This protein has broad tissue distribution, and is reduced in expression on Rh erythrocytes.