



Recombinant Protein Technical Manual

Recombinant Rat Interferon γ /IFNG Protein (Fc Tag)(Active)

RPES3869

Product Data:

Product SKU: RPES3869

Size: 20 μ g

Species: Rat

Expression host: HEK293 Cells

Uniprot: P01581

Protein Information:

Molecular Mass: 42.5 kDa

AP Molecular Mass: 52 kDa

Tag: C-Fc

Bio-activity: 1. Measured in antiviral assay using L929 cells infected with vesicular stomatitisvirus (VSV). The ED50 for this effect is typically 1-5 ng/mL. 2. Measured by its binding ability in a functional ELISA. Immobilized rat IFNG-Fc at 10 μ g/ml (100 μ l/well) can bind biotinylated rat IFNGR-Fc. The EC50 of biotinylated rat IFNGR-Fc is 20.1-46.9 ng/ml.

Purity: > 92 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per μ g of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: IFNG2

Immunogen Information:

Sequence: Val 117-Ser 268

Background:

IFN gamma, also known as IFNG, is a secreted protein which belongs to the type I I interferon family. IFN gamma is produced predominantly by natural killer and natural killer T cells as part of the innate immune response, and by CD4 and CD8 cytotoxic T lymphocyte effector T cells once antigen-specific immunity develops. IFN gamma has antiviral, immunoregulatory, and anti-tumor properties. IFNG, in addition to having antiviral activity, has important immunoregulatory functions, it is a potent activator of macrophages, and has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons. The IFNG monomer consists of a core of six α -helices and an extended unfolded sequence in the C-terminal region. IFN gamma is critical for innate and adaptive immunity against viral and intracellular bacterial infections and for tumor control. Aberrant IFN gamma expression is associated with a number of autoinflammatory and autoimmune diseases. The importance of IFN gamma in the immune system stems in part from its ability to inhibit viral replication directly, and most importantly from its immunostimulatory and immunomodulatory effects. IFNG also promotes NK cell activity.