



Recombinant Protein Technical Manual
Recombinant Human TrkA/NTRK1 Protein (aa 285-413, His Tag)
RPES3903

Product Data:

Product SKU: RPES3903

Size: 50µg

Species: Human

Expression host: E. coli

Uniprot: NP_002520.2

Protein Information:

Molecular Mass: 15.1 kDa

AP Molecular Mass: 16 kDa

Tag: N-His

Bio-activity:

Purity: > 97 % as determined by reducing SDS-PAGE.

Endotoxin: Please contact us for more information.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile 50mM Tris, 200mM NaCl, pH 8.0

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: MTC;p140-TrkA;TRK;Trk-A;TRK1;TRKA

Immunogen Information:

Sequence: Pro 285-Glu 413

Background:

TRKA is a member of the neurotrophic tyrosine kinase receptor (NTRK) family. It is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Isoform TrkA-III promotes angiogenesis and has oncogenic activity when overexpressed. Isoform TrkA-I is found in most non-neuronal tissues. Isoform TrkA-II is primarily expressed in neuronal cells. TrkA-III is specifically expressed by pluripotent neural stem and neural crest progenitors. The presence of NTRK1 leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in TRKA gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. It was originally identified as an oncogene as it is commonly mutated in cancers, particularly colon and thyroid carcinomas. TRKA is required for high-affinity binding to nerve growth factor (NGF), neurotrophin-3 and neurotrophin-4/5 but not brain-derived neurotrophic factor (BDNF). Known substrates for the Trk receptors are SHC1, PI 3-kinase, and PLC-gamma. NTRK1 has a crucial role in the development and function of the nociceptive reception system as well as establishment of thermal regulation via sweating. It also activates ERK1 by either SHC1- or PLC-gamma-dependent signaling pathway. Defects in NTRK1 are a cause of congenital insensitivity to pain with anhidrosis and thyroid papillary carcinoma.