



Recombinant Protein Technical Manual
Recombinant Mouse Arylsulfatase A/ARSA Protein
(His Tag)(Active)
RPES4162

Product Data:

Product SKU: RPES4162

Size: 20µg

Species: Mouse

Expression host: HEK293 Cells

Uniprot: NP_033843.2

Protein Information:

Molecular Mass: 53.5 kDa

AP Molecular Mass: 60 kDa

Tag: C-His

Bio-activity: Measured by its ability to cleave p-Nitrocatechol Sulfate (PNCS). The specific activity is >100 pmoles/min/µg.

Purity: > 97 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile 25mM Tris, 0.15mM NaCl, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: As-2;AS-A;As2;ASA;AW212749;TISP73

Immunogen Information:

Sequence: Met 1-Ser 506

Background:

Arylsulfatase A (ARSA) is synthesized as a 52KDa lysosomal enzyme. It is a member of the sulfatase family that is required for the lysosomal degradation of cerebroside-3-sulfate, a sphingolipid sulfate ester and a major constituent of the myelin sheath. Arylsulfatase A is activated by a required co- or posttranslational modification with the oxidation of cysteine to formylglycine. Metachromatic leukodystrophy (MLD) is a lysosomal storage disease in the central and peripheral nervous systems with severe and progressive neurological symptoms caused by the deficiency of Arylsulfatase A. Deficiency of this enzyme is also found in apparently healthy individuals, a condition for which the term pseudodeficiency is introduced. ARSA forms dimers after receiving three N-linked oligosaccharides in the endoplasmic reticulum, and then the dimers are transported to the Golgi where they receive mannose 6-phosphate recognition markers. And thus, ARSA is transported and delivered to dense lysosomes in a mannose 6-phosphate receptor-dependent manner. It has been shown that within the lysosomes, the ARSA dimers can oligomerize to an octamer in a pH-dependent manner. The ARSA deficiency leads to metachromatic leukodystrophy (MLD), a lysosomal storage disorder associated with severe and progressive demyelination in the central and peripheral nervous system. Additionally, the serum level of arylsulfatase A might be helpful in diagnosis of lung and central nervous system cancer.