

Recombinant Protein Technical Manual Recombinant Human TROP2/TACSTD2 Protein (aa 1-

**274, His Tag)** RPES4204

## Product Data:

Product SKU: RPES4204

Species: Human

**Size:** 100µg

Expression host: HEK293 Cells

**Uniprot:** NP\_002344.2

Drotoin	<b>Information:</b>
Protein	

Molecular Mass:	29.4 kDa
AP Molecular Mass:	42-48 kDa
Tag:	C-His
Bio-activity:	
Purity:	> 96 % as determined by reducing SDS-PAGE.
Endotoxin:	< 1.0 EU per $\mu g$ as determined by the LAL method.
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	EGP;EGP1;GA733;GA7331;GP50;M1S1;TACD2;TROP-2;TROP2

## Sequence: Met 1-Thr 274

## Background:

TROP-2, also referred to as tumor associated calcium signal transducer 2 (TACSTD2), GA733 or M1S1, is a cell surface glycoprotein highly expressed in a wide variety of epithelial cancers. In contrast, there is little or no expression of Trop-2 in adult somatic tissue. Because it is a cell surface protein that is selectively expressed in tumor cells, Trop-2 is a potential therapeutic target. The cytoplasmic tail of Trop-2 possesses potential serine and tyrosine phosphorylation sites and a phosphatidyl-inositol binding consensus sequence. Trop-2 transduces an intracellular calcium signal, are consistent with the hypothesis that it acts as a cell surface receptor and support a search for a physiological ligand. TROP2 encoding by an intronless gene was originally defined by the monoclonal antibody GA733, and is a member of a family of at least two type I membrane proteins. The other known member is GA733-2, also called EpCAM and TROP1. It has been suggested by studies that the GA733 gene was formed by the retroposition of the GA733-2 gene via an mRNA intermediate.