



Recombinant Protein Technical Manual

Recombinant Rat CD99/MIC2 Protein (Fc Tag)

RPES4394

Product Data:

Product SKU: RPES4394

Size: 50µg

Species: Rat

Expression host: HEK293 Cells

Uniprot: B4F7A5

Protein Information:

Molecular Mass: 37.6 kDa

AP Molecular Mass: 47 kDa

Tag: C-Fc

Bio-activity:

Purity: > 95 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: CD99

Immunogen Information:

Sequence: Met1-Gly127

Background:

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD99 is a transmembrane protein expressed on most hematopoietic cells, endothelial cells and at the borders between confluent cells. CD99 is also found expressed in the development of normal ovary and testis as well as in 25 sex cord-stromal tumors, 7 epithelial neoplasms, and 6 germ cell tumors. CD99 may be a useful marker for sex cord-stromal tumors and that its degree of reactivity correlates with the degree of differentiation in Sertoli-Leydig cell tumors. Additionally, CD99 might aid in distinguishing granulosa cell tumors of the ovary from poorly differentiated carcinomas and it has been reported to be a sensitive and specific marker for Ewing's sarcoma and primitive neuroectodermal tumor.