

Recombinant Protein Technical Manual Recombinant Rat CLEC4A3 Protein (Fc Tag)

RPES4426

## Product Data:

Product SKU: RPES4426	<b>Size:</b> 20µg
Species: Rat	Expression host: HEK293 Cells

Uniprot: Q5YIS0

## **Protein Information:**

Molecular Mass:	48.3 kDa
worecular wass:	48.3 KDa
AP Molecular Mass:	58 kDa
Tag:	N-Fc
Bio-activity:	
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU per $\mu g$ of the protein as determined by the LAL method
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	CLEC4A3

## Sequence: Leu68-Leu237

## Background:

CLEC4A3 contains 1 C-type lectin domain and belongs to the C-type lectin-like domain-containing (CLEC) family. Lectins are proteins that are able to recognize and bind with specific carbohydrate molecules. C-type lectins are an important group of proteins found in the immune system of animals. These lectins are named C-type because of their calcium dependent carbohydrate recognition domain (CRD). In the immune system, C-type lectins act as recognition molecules by binding to foreign microorganisms. They also promote the movement and selective adhesion of white blood cells. The C-type lectin has a three-dimensional fold, the CRD, in which calcium ions contribute to the lectin's ability to recognize and bind carbohydrates. In the immune system, carbohydrate recognition contributes to the ability of immune cells to move from one area of the body to another. It also allows immune cells to identify and discriminate between proteins that belong to the host and those that belong to foreign organisms. There are a number of different C-type lectin subfamilies, including collectins, selectins, proteoglycans, and lymphocyte lectins.