



Recombinant Protein Technical Manual
Recombinant Human Complexin-2/CPLX2 Protein
(His Tag)
RPES4707

Product Data:

Product SKU: RPES4707

Size: 20µg

Species: Human

Expression host: E. coli

Uniprot: Q6PUV4

Protein Information:

Molecular Mass: 16.8 kDa

AP Molecular Mass: 21 kDa

Tag: N-His

Bio-activity:

Purity: > 90 % as determined by reducing SDS-PAGE.

Endotoxin: Please contact us for more information.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: 921-L;CPX-2;CPX2;Hfb1

Immunogen Information:

Sequence: Asp 2-Lys 134

Background:

Complexin-2 (CPLX2), a member of the complexin/synaphin family, is a soluble pre-synaptic protein believed to regulate neurotransmitter release from pre-synaptic terminals. Complexins are soluble proteins that regulate the activity of soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complexes necessary for vesicle fusion. Complexins are unable to bind to monomeric SNARE proteins but bind with high affinity to ternary SNARE complexes and with lower affinity to target SNARE complexes. Complexin 1 (CX1) and complexin 2 (CX2) are presynaptic proteins that modulate neurotransmitter release and are used as markers of inhibitory and excitatory synapses, respectively. CPLX2 is localized in pre-synaptic terminals in mature brain. The G71-P89 region of CPLX2 is essential and sufficient for preferential axonal distribution. CPLX2 participates in the Ca(2+)-sensitive regulatory pathway for zymogen granule exocytosis. Complexin-2 is a key player in normal neurological function, and its downregulation could lead to changes in neurotransmitter release sufficient to cause significant behavioural abnormalities such as depression. It is involved in synaptogenesis and the modulation of neurotransmitter release.