



Recombinant Protein Technical Manual
Recombinant Rat CD10/Nepriylsin Protein (His Tag)
RPES4710

Product Data:

Product SKU: RPES4710

Size: 10µg

Species: Rat

Expression host: HEK293 Cells

Uniprot: P07861

Protein Information:

Molecular Mass: 82.5 kDa

AP Molecular Mass: 93 kDa

Tag: N-His

Bio-activity:

Purity: > 95 % as determined by SDS-PAGE

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: MME

Immunogen Information:

Sequence: Tyr52-Trp750

Background:

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 10 (CD10), also known as Neprilysin and neutral endopeptidase, is a member of the CD system. CD10 is a zinc-dependent metalloprotease enzyme that had function to degrade a number of small secreted peptides such as the amyloid beta peptide. It exist as a membrane-bound protein and have high concentration in kidney and lung tissues. Mutations in the CD10 gene can induce the familial forms of Alzheimer's disease, providing strong evidence for the protein's association with the Alzheimer's disease process. CD10 is also associated with other biochemical processes.