

Recombinant Protein Technical Manual Recombinant Rat CD63/Tspan-30 Protein (Fc Tag) RPES4719

**Product Data:** 

Product SKU: RPES4719	<b>Size:</b> 20µg
Species: Rat	Expression host: HEK293 Cells

Uniprot: P28648

Drot	an	Intor	mation	
	GILL			

Molecular Mass:	38.1 kDa
AP Molecular Mass:	42-47 kDa
Tag:	N-Fc
Bio-activity:	
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin:	< 1.0 EU per $\mu g$ of the protein as determined by the LAL method
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	CD63

## Sequence: Ala103-Val203

## **Background:**

The cluster of differentiation (CD) system is commonly used as cell markers in immunophynotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. Cluster of differentiation 63 (CD63) is a member of the CD family and the transmembrane 4 superfamily, also known as the tetraspanin family. CD63 is a cellular surface glycoprotein characterized by the presence of four bydrophobic domains. CD63 had functions in mediating signal transduction processes and then regulate variety of cellular processes such as cell proliferation, activation and motility. It has reported that CD63 protein associated with tumor progression and served as a blood platlet activation marker and the deficiency of this protein may be associated with Hermansky-Pudlak syndrome.