



Recombinant Protein Technical Manual  
Recombinant Mouse CD28/TP44 Protein (His Tag)  
RPES4735

#### Product Data:

**Product SKU:** RPES4735

**Size:** 100µg

**Species:** Mouse

**Expression host:** HEK293 Cells

**Uniprot:** NP\_031668.3

#### Protein Information:

**Molecular Mass:** 16.5 kDa

**AP Molecular Mass:**

**Tag:** C-His

**Bio-activity:**

**Purity:** > 95 % as determined by SDS-PAGE

**Endotoxin:** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from sterile PBS, pH 7.4

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:**

**Synonyms:** T-cell-specific surface glycoprotein CD28;CD28;CH2989K17.3

## Immunogen Information:

**Sequence:** Met 1-Lys 149

## Background:

CD28(Cluster of Differentiation 28) is a disulphide-bonded glycoprotein belonging to the immunoglobulin (Ig) superfamily, and structurally consists of a single Ig V-like extracellular domain, a transmembrane domain and an intracellular domain. Mouse CD28 is constitutively expressed on the surface of all murine T cells and on developing thymocytes as disulfide-linked homodimers or as monomers. CD28 can binds the B7 and B7-2 ligand, and together perform important functions in the T and B cell response pathways. B7/CD28 family members, which can augment or antagonize T-cell receptor signaling, in the regulation of central and peripheral T-cell tolerance. CD28 is thus involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival.