



Recombinant Protein Technical Manual
Recombinant Human Ficolin/Ficolin-A/FCN1 Protein
(His Tag)(Active)
RPES5059

Product Data:

Product SKU: RPES5059

Size: 50µg

Species: Human

Expression host: HEK293 Cells

Uniprot: NP_001994.2

Protein Information:

Molecular Mass: 33.6 kDa

AP Molecular Mass: 33.6 kDa

Tag: C-His

Bio-activity: Measured by its binding ability in a functional ELISA.1. Immobilized FCN1 at 10 µg/ml can bind biotinylated recombinant human Fetuin-A with a linear range of 16-2000 ng/ml.2. Immobilized FCN1 at 10 µg/ml can bind biotinylated recombinant mouse Fetuin-A with a linear range of 16-2000 ng/ml.

Purity: > 95 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: FCN1;FCNA;FCNM

Immunogen Information:

Sequence: Met 1-Ala 326

Background:

Ficolins are humoral molecules of the innate immune systems which recognize carbohydrate molecules on pathogens, apoptotic and necrotic cells. The Ficolin family of proteins are characterized by the presence of a leader peptide, a short N-terminal segment, followed by a collagen-like region, and a C-terminal fibrinogen-like domain. Ficolins are humoral molecules of the innate immune systems which recognize carbohydrate molecules on pathogens, apoptotic and necrotic cells. Three Ficolins have been identified in humans: L-Ficolin, H-Ficolin and M-Ficolin (also referred to as Ficolin-2, -3 and , respectively). They are soluble oligomeric defence proteins with lectin-like activity and they are structurally similar to the human collectins, mannan-binding lectin (MBL) and surfactant protein A and D. Dysfunction or abnormal expressions of Ficolins may involved in the pathogenesis of human diseases including infectious and inflammatory diseases, autoimmune disease and clinical syndrome of preeclampsia. They are soluble oligomeric defence proteins with lectin-like activity and they are structurally similar to the human collectins, mannan-binding lectin (MBL) and surfactant protein A and D. Upon recognition of the infectious agent, the Ficolins act through two distinct routes: initiate the lectin pathway of complement activation through attached serine proteases (MASPs), and a primitive opsonophagocytosis thus limiting the infection and concurrently orchestrating the subsequent adaptive clonal immune response. Ficolin (FCN1) is predominantly expressed in the peripheral blood leukocytes.