



Recombinant Protein Technical Manual

Recombinant Human EphB6 Protein (His Tag)(Active)
RPES5142

Product Data:

Product SKU: RPES5142

Size: 100µg

Species: Human

Expression host: HEK293 Cells

Uniprot: NP_004436.1

Protein Information:

Molecular Mass: 61.6 kDa

AP Molecular Mass: 60-70 kDa

Tag:

Bio-activity: 1. Measured by its binding ability in a functional ELISA.2. Immobilized recombinant human EphB6 at 10 µg/ml (100 µl/well) can bind human EphrinB1 with a linear range of 32-800 ng/ml.3. Immobilized recombinant human EphB6 at 10 µg/ml (100 µl/well) can bind human EphrinB2 with a linear range of 1.28-32 ng/ml.

Purity: > 92 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per µg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: HEP

Immunogen Information:

Sequence: Met 1-Ser 579

Background:

Ephrins are divided into the ephrin-A (EFNA) class and the ephrin-B (EFNB) class based on their structures and sequence relationships. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. EphB6 is an unusual Eph receptor, lacking catalytic capacity due to alterations in its kinase domain. Interestingly, increased metastatic activity is associated with reduced EphB6 receptor expression in several tumor types, including breast cancer. This emphasizes the potential of EphB6 to act as a suppressor of cancer aggressiveness. EphB6 suppress cancer invasiveness through c-Cbl-dependent signaling, morphologic changes, and cell attachment and indicate that EphB6 may represent a useful prognostic marker and a promising target for therapeutic approaches. EphB6 can both positively and negatively regulate cell adhesion and migration, and suggest that tyrosine phosphorylation of the receptor by an Src family kinase acts as the molecular switch for the functional transition. In addition, Ephrin-B2 may be a physiological ligand for the EphB6 receptor.