

Recombinant Protein Technical Manual Recombinant Human PD-L2/CD273 Protein (His & Avi Tag)

Product Data:

Product SKU: RPES5153 **Size:** 20μg

Species: Human Cells

RPES5153

Uniprot: Q9BQ51

Protein Information:

Molecular Mass: 25.4 kDa

AP Molecular Mass: 38-60 kDa

Tag: C-His-Avi

Bio-activity:

Purity: > 95% as determined by reducing SDS-PAGE.

Endotoxin: $< 1.0 \text{ EU per } \mu\text{g}$ as determined by the LAL method.

Storage: Lyophilized protein should be stored at < -20°C, though stable at room

temperature for 3 weeks. Reconstituted protein solution can be stored at $4-7^{\circ}$ C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.

Reconstitution: Please refer to the printed manual for detailed information.

Application:

Synonyms: Programmed Cell Death 1 Ligand 2; PD Ligand 2; PD-L2; PDCD1 Ligand

2; Programmed Death Ligand 2; Butyrophilin B7-DC; B7-DC; CD273; PDCD1LG2;

B7DC;CD273; PDCD1L2; PDL2

Immunogen Information:

Sequence: Leu20-Pro219

Background:

Programmed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG family. PDCD1LG2 contains one Ig-like C2-type domain and one Ig-like V-type domain. PDCD1LG2 is highly expressed in the heart, placenta, pancreas, lung and liver; it is weakly expressed in the spleen, lymph nodes, and thymus. PDCD1LG2 is involved in the costimulatory signal, essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.