

# Recombinant Protein Technical Manual Recombinant Human CD44 Protein (His Tag)

**RPES5202** 

#### Product Data:

**Product SKU:** RPES5202 **Size:** 10μg

Species: Human Cells

**Uniprot:** P16070

### **Protein Information:**

Molecular Mass: 23.1 kDa

AP Molecular Mass: 44 kDa

**Tag:** C-6His

**Bio-activity:** 

**Purity:** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin:**  $< 1.0 \text{ EU per } \mu\text{g}$  as determined by the LAL method.

**Storage:** Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:** 

**Synonyms:** CDW44;CSPG8;ECMR-III;HCELL;HUTCH-I;IN;LHR;MC56;MDU2;MDU3;MIC4;Pgp1;

Epican; Extracellular Matrix Receptor III; ECMR-III; GP90 Lymphocyte

Homing/Adhesion Receptor; HUTCH-I; Heparan Sulfate Proteoglycan; Hermes Antigen; Hyaluronate Receptor; Phagocytic Glycoprotein 1; PGP; Phagocytic

Glycoprotein I; PGP-I; CD44

## Immunogen Information:

Sequence: Gln21-Pro220

## **Background:**

CD44 is a cell-surface receptor for hyaluronic acid and also interacts with other ligands, such as osteopontin, collagens, and matrix metalloproteinases. A large number of CD44 isoforms can be generated by the insertion of different combinations of at least nine exons. Increased CD44 antigen is associated with relapses in non-small cell lung cancers. Furthermore, an increasing quantity of evidence suggests that CD44 has various functions related to inflammatory disease. CD44 deficiency induces severe liver injury. CD44-hyaluronate mediates in lymphocyte T and monocyte adhesion to the endothelium, stimulates proinflammatory cytokine release from macrophages and participates in dedifferentiation phenotype of smooth muscle cells from contractile state to synthetic one.