# Feline SDF 1b Recombinant Protein

## **RPPB5746**



| Product Information | Protein Information   |
|---------------------|---|
| Product SKU:        | Protein description:  |
| RPPB5746            | Stromal Cell-Derived Factor-1 beta Feline Recombinant produced in E.Coli is a non-glycosylated,       |
|                     | Polypeptide chain containing 72 amino acids and having a molecular mass of 8526 Dalton. The SDF-1b is |
| Accession:          | purified by proprietary chromatographic techniques.   |
| O62657              |   |
|                     | Appearance:   |
| Host:               | Sterile Filtered White lyophilized (freeze-dried) powder.   |
| Escherichia Coli.   |   |
|                     | Synonyms:   |
|                     | SDF-1, CXCL12, Pre-B cell growth-stimulating factor, PBSF, hIRH, chemokine (C-X-C motif) ligand 12,   |
|                     | SDF1, SDF1B, TPAR1, SCYB12, SDF-1b, TLSF-b, 12-O-tetradecanoylphorbol 13-acetate repressed protein    |
|                     | 1, Thymic lymphoma cell-stimulating factor, TLSF.   |
|                     | Formulation:  |
|                     | The protein was lyophilized from a concentrated (1 mg/ml) sterile solution containing no additives.   |

#### Purity:

Greater than 97.0% as determined by(a) Analysis by RP-HPLC.(b) Analysis by SDS-PAGE.

#### Solubility:

It is recommended to reconstitute the lyophilized SDF-1b in sterile  $18M\Omega$ -cm H2O not less than  $100\mu$ g/ml, which can then be further diluted to other aqueous solutions.

#### Stability:

Lyophilized SDF-1b although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution CXCL12 should be stored at 4°C between 2-7 days and for future use below - 18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Please prevent freeze-thaw cycles.

#### **Amino Acid Sequence:**

KPVSLSYRCP CRFFESHVAR ANVKHLKILN TPNCALQIVA RLKNNNRQVC IDPKLKWIQE YLEKALNKRF KM.

### **Biological Activity:**

The specific activity as determined by its ability to chemoattract human peripheral T cells stimulated with IL-2 and PHA in a concentration range from 10-75 ng/ml corresponding to a Specific Activity of 13,334-100,000IU/mg.