

CARP01337

Product Information

Species:

SARS-COV-2

Gene ID

43740568
(RBD) (L452R, E484Q)

Category:

Recombinant Protein

Swiss Prot:

PODTC2

Tags:

C-6xHis

Synonyms:

S1-RBD protein;NCP-CoV RBD Protein;novel coronavirus RBD Protein;2019-nCoV RBD Protein;S glycoprotein Subunit1 RBD Protein

Antibody Information

Purification:

>97% by SDS-PAGE

Source:

HEK293 cells

Endotoxin:

< 0.1 EU/μg of the protein by LAL method

Formulation:

Supplied as a 0.22 μm filtered solution in PBS, pH 7.4

Reconstitution:

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water

Background

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate

Description

SARS-CoV-2 Delta Variant B.1.617.2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Arg319-Phe541(L452R,E484Q)) of SARS-COV-2 RBD (Accession #YP_009724390.1) fused with a 6xHis tag at the C-terminus.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Product Images



Recombinant SARS-COV-2 Spike RBD(L452R,E484Q) Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 36kDa.