

FITC Anti-Mouse/Human CD11b Antibody [M1/70]

AGEL0306

Description

This FITC Anti-Mouse/Human CD11b Antibody [M1/70] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL0306
Contents:	100 Tests, 200 Tests, 50 Tests Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	M1/70
Synonyms:	CD11 antigen-like family member B, CD11b, CR-3 alpha chain, Integrin alpha-M, Itgam, Leukocyte adhesion receptor MO1
Applications:	FCM
Reactivity:	Human, Mouse
Immunogen:	-

Antibody Data

Uniprot ID:	P05555, P11215
Gene ID:	16409, 3684
Swissprot:	P05555P11215
Host Species:	Rat
Isotype:	Rat IgG2b, κ
Isotype Control:	FITC Rat IgG2b, κ Isotype Control[LTF-2]

Conjugation:	FITC
Conjugation Information:	FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
Purification:	-
Target:	CD11b
Cellular Localization:	Membrane
Tissue Specificity:	-
Verified Samples:	-
Concentration:	5 µL/Test

Preparation & Storage

Storage: This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.

Shipping: Ice bag

Recommended Dilution: -

Recommended Usage:

Application	Recommended Usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.