

# Purified Anti-Mouse CD3epsilon Antibody [145-2C11]

AGEL0374

## Description

---

This Purified Anti-Mouse CD3epsilon Antibody [145-2C11] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	AGEL0374
<b>Contents:</b>	100µg, 25µg Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Monoclonal Antibody
<b>Clonality:</b>	Monoclonal
<b>Clone:</b>	145-2C11
<b>Synonyms:</b>	CD3E, CD3e, T-cell surface antigen T3/Leu-4 epsilon chain, T-cell surface glycoprotein CD3 epsilon chain, T3E
<b>Applications:</b>	<a href="#">FCM</a>
<b>Reactivity:</b>	Mouse
<b>Immunogen:</b>	Recombinant Mouse CD3ε protein

## Antibody Data

---

<b>Uniprot ID:</b>	-
<b>Gene ID:</b>	-
<b>Swissprot:</b>	P22646
<b>Host Species:</b>	Armenian Hamster
<b>Isotype:</b>	Armenian Hamster IgG
<b>Isotype Control:</b>	-

**Manufacturers Statement:** This final kit system is assembled and quality-released by Assay Genie Limited.

<b>Conjugation:</b>	-
<b>Conjugation Information:</b>	-
<b>Buffer:</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.
<b>Purification:</b>	>98%, Protein A/G purified
<b>Target:</b>	CD3εμ
<b>Cellular Localization:</b>	-
<b>Tissue Specificity:</b>	-
<b>Verified Samples:</b>	Verified Samples in FCM: C57/BL6 Mouse splenocytes
<b>Concentration:</b>	≥ 1 mg/mL

## Preparation & Storage

---

**Storage:** Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.

**Shipping:** Ice bag

**Recommended Dilution:** FCM 2 μg/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

**Recommended Usage:**

Application	Recommended Usage
-	-

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.