

# Anti-Human/Monkey CD154/CD40L Antibody [5C8] -Low Endotoxin Purified

AGEL1603

## Description

---

This Anti-Human/Monkey CD154/CD40L Antibody [5C8] -Low Endotoxin Purified is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	AGEL1603
<b>Contents:</b>	100µg, 25µg Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Monoclonal Antibody
<b>Clonality:</b>	Monoclonal
<b>Clone:</b>	5C8
<b>Synonyms:</b>	T-BAM, TNFSF5, TRAP, gp39, CD40L
<b>Applications:</b>	<a href="#">FCM</a>
<b>Reactivity:</b>	Human, Monkey
<b>Immunogen:</b>	Recombinant Human CD154 protein

## Antibody Data

---

<b>Uniprot ID:</b>	-
<b>Gene ID:</b>	-
<b>Swissprot:</b>	P29965
<b>Host Species:</b>	Mouse
<b>Isotype:</b>	Mouse IgG2a, κ
<b>Isotype Control:</b>	-

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

<b>Conjugation:</b>	-
<b>Conjugation Information:</b>	-
<b>Buffer:</b>	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.
<b>Purification:</b>	>98%, Protein A/G purified
<b>Target:</b>	CD154
<b>Cellular Localization:</b>	Secreted, membrane
<b>Tissue Specificity:</b>	-
<b>Verified Samples:</b>	Verified Samples in FCM: HEK293T cells transfected with pcDNA3.1 plasmid encoding Human CD154 gene
<b>Concentration:</b>	≥ 1 mg/mL

## Preparation & Storage

---

**Storage:** Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.

**Shipping:** Ice bag

**Recommended Dilution:** FCM 2 µg/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

**Recommended Usage:**

Application	Recommended Usage
-	-

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.