

# Anti-Mouse IL-6 Antibody [MP5-20F3] -Low Endotoxin Purified

AGEL2070

## Description

---

This Anti-Mouse IL-6 Antibody [MP5-20F3] -Low Endotoxin Purified is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	AGEL2070
<b>Contents:</b>	100µg, 25µg Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Monoclonal Antibody
<b>Clonality:</b>	Monoclonal
<b>Clone:</b>	MP5-20F3
<b>Synonyms:</b>	BSF, Interleukin, Interleukin HP, IFNB, B-cell hybridoma growth factor, BSF2, CDF, HGF, HSF, IFNB2, IL 6, Interleukin HP-1, Interleukin-6, IL-6, IL6
<b>Applications:</b>	<a href="#">FCM</a>
<b>Reactivity:</b>	Mouse
<b>Immunogen:</b>	Recombinant Human IL-6 protein

## Antibody Data

---

<b>Uniprot ID:</b>	-
<b>Gene ID:</b>	-
<b>Swissprot:</b>	P08505
<b>Host Species:</b>	Rat
<b>Isotype:</b>	Rat IgG1, κ

<b>Isotype Control:</b>	-
<b>Conjugation:</b>	-
<b>Conjugation Information:</b>	-
<b>Buffer:</b>	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.
<b>Purification:</b>	-
<b>Target:</b>	IL-6
<b>Cellular Localization:</b>	Secreted
<b>Tissue Specificity:</b>	-
<b>Verified Samples:</b>	Verified Samples in FCM: HEK293T cells transfected with pcDNA3.1 plasmid encoding Mouse IL-6 gene
<b>Concentration:</b>	≥ 1 mg/mL

## Preparation & Storage

---

**Storage:** Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.

**Shipping:** Ice bag

**Recommended Dilution:** FCM 2 µg/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

**Recommended Usage:**

Application	Recommended Usage
-	-

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.