

# PE/GenieFluor 594 Anti-Human CD38 Antibody [HIT2]

AGEL3146

## Description

---

This PE/GenieFluor 594 Anti-Human CD38 Antibody [HIT2] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	AGEL3146
<b>Contents:</b>	100 Tests, 20 Tests, 200 Tests Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Monoclonal Antibody
<b>Clonality:</b>	Monoclonal
<b>Clone:</b>	HIT2
<b>Synonyms:</b>	2'-phospho-ADP-ribosyl cyclase, 2'-phospho-cyclic-ADP-ribose transferase, ADP-ribosyl cyclase 1, ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1, ADPRC 1, CD38, Cyclic ADP-ribose hydrolase 1, T10, cADPr hydrolase 1
<b>Applications:</b>	<b>FCM</b>
<b>Reactivity:</b>	Human
<b>Immunogen:</b>	-

## Antibody Data

---

<b>Uniprot ID:</b>	P28907
<b>Gene ID:</b>	952
<b>Swissprot:</b>	P28907
<b>Host Species:</b>	Mouse
<b>Isotype:</b>	Mouse IgG1, κ

**Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.**

<b>Isotype Control:</b>	PE/GenieFluor 594 Mouse IgG1, κ Isotype Control[MOPC-21]
<b>Conjugation:</b>	PE/GenieFluor594
<b>Conjugation Information:</b>	PE/GenieFluor 594 is designed to be excited by the blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 620 nm (e.g., a 610/20 nm bandpass filter).
<b>Buffer:</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
<b>Purification:</b>	-
<b>Target:</b>	CD38
<b>Cellular Localization:</b>	Membrane
<b>Tissue Specificity:</b>	-
<b>Verified Samples:</b>	-
<b>Concentration:</b>	5 µL/Test

## Preparation & Storage

---

<b>Storage:</b>	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping:</b>	Ice bag
<b>Recommended Dilution:</b>	-

<b>Recommended Usage:</b>	<b>Application</b>	<b>Recommended Usage</b>
	FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:**

Centrifuge before opening to ensure complete recovery of vial contents.