

GenieFluor 647 Anti-Mouse Perforin Antibody [S16009A]

AGEL3309

Description

This GenieFluor 647 Anti-Mouse Perforin Antibody [S16009A] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL3309
Contents:	100 Tests, 200 Tests, 50 Tests Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	S16009A
Synonyms:	PfnPfn-1Pfp, Cytolysin, Lymphocyte pore-forming protein, Prf1
Applications:	ICFCM
Reactivity:	Mouse
Immunogen:	-

Antibody Data

Uniprot ID:	P10820
Gene ID:	18646
Swissprot:	P10820
Host Species:	Rat
Isotype:	Rat IgG2a, κ
Isotype Control:	GenieFluor 647 Rat IgG2a, κ Isotype Control[2A3]

Conjugation:	GenieFluor647
Conjugation Information:	GenieFluor 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter).
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
Purification:	-
Target:	Perforin
Cellular Localization:	Cytoplasm
Tissue Specificity:	-
Verified Samples:	-
Concentration:	5 µL/Test

Preparation & Storage

Storage:	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping:	Ice bag
Recommended Dilution:	-

Recommended Usage:

Application	Recommended Usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use

Protein Quantification (Optional):

To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.