

GenieFluor 647 Anti-Mouse CD150 Antibody [TC15-12F12.2]

AGEL3371

Description

This GenieFluor 647 Anti-Mouse CD150 Antibody [TC15-12F12.2] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

| | |
|----------------------|---|
| SKU: | AGEL3371 |
| Contents: | 100µg, 25µg Bradford Reagent: 1 vial (2ml) |
| Category: | Monoclonal Antibody |
| Clonality: | Monoclonal |
| Clone: | TC15-12F12.2 |
| Synonyms: | CD 150, Signaling lymphocytic activation molecule, Slam, Slamf1 |
| Applications: | FCM |
| Reactivity: | Mouse |
| Immunogen: | - |

Antibody Data

| | |
|-------------------------|--|
| Uniprot ID: | Q9QUM4 |
| Gene ID: | 27218 |
| Swissprot: | Q9QUM4 |
| Host Species: | Rat |
| Isotype: | Rat IgG2a, λ |
| Isotype Control: | GenieFluor 647 Rat IgG2a, λ Isotype Control[B39-4] |

| | |
|---------------------------------|---|
| Conjugation: | GenieFluor647 |
| Conjugation Information: | GenieFluor 647 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 670 nm (e.g., a 660/20 nm bandpass filter). |
| Buffer: | Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. |
| Purification: | - |
| Target: | CD150/SLAM |
| Cellular Localization: | Membrane |
| Tissue Specificity: | - |
| Verified Samples: | - |
| Concentration: | 0.5 mg/mL |

Preparation & Storage

| | |
|------------------------------|---|
| Storage: | This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. |
| Shipping: | Ice bag |
| Recommended Dilution: | - |

| Recommended Usage: | Application | Recommended Usage |
|---------------------------|-------------|---|
| | FCM | Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10 ⁶ cells in 100 µL volume] |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.