

GenieFluor 700 Anti-Mouse Ly-6G/Ly-6C (Gr-1) Antibody [RB6-8C5]

AGEL4146

Description

This GenieFluor 700 Anti-Mouse Ly-6G/Ly-6C (Gr-1) Antibody [RB6-8C5] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL4146
Contents:	100µg, 25µg Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	RB6-8C5
Synonyms:	Gr-1, Gr1, Ly-6G/Ly-6C, Ly6G/Ly6C
Applications:	FCM
Reactivity:	Mouse
Immunogen:	-

Antibody Data

Uniprot ID:	P35461, P0CW03,
Gene ID:	546644, 17067
Swissprot:	P35461P0CW03
Host Species:	Rat
Isotype:	Rat IgG2b, κ
Isotype Control:	GenieFluor 700 Rat IgG2b, κ Isotype Control[LTF-2]

Conjugation:	GenieFluor700
Conjugation Information:	GenieFluor 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
Buffer:	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.
Purification:	-
Target:	Ly-6G/Ly-6C (Gr-1)
Cellular Localization:	Membrane
Tissue Specificity:	-
Verified Samples:	-
Concentration:	-

Preparation & Storage

Storage:	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping:	Ice bag
Recommended Dilution:	-

Recommended Usage:	Application	Recommended Usage
	FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/106 cells in 100 µL volume]

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.