

Anti-Human CD62L Antibody [DREG56] -Low Endotoxin, Azide Free

AGEL4985

Description

This Anti-Human CD62L Antibody [DREG56] -Low Endotoxin, Azide Free is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL4985
Contents:	25µg, 100µg, 1mg Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	DREG56
Synonyms:	Leukocyte adhesion molecule, Leukocyte surface antigen Leu, Leukocyte-endothelial cell adhesion molecule, LECAM, LAM, LEU, SELL, CD62L, LAM1, LECAM1, LEU8, LNHR, LSEL, LYAM1, PLNHR, TQ1, CD62 antigen-like family member L, CD62antigen-likefamilymemberL, gp90-MEL, LAM-1, Leu-8, Leukocyte adhesion molecule 1, Leukocyte surface antigen Leu-8, Leukocyte-endothelial cell adhesion molecule 1, L-selectin, Ly22, Ly-22, Lyam-1, Lymph node homing receptor, Lymphnodehomingreceptor, 6&, aposulfated Lewis X-X, LECAM-1, LAM-1
Applications:	FCM
Reactivity:	Human
Immunogen:	Recombinant Human CD62L protein

Antibody Data

Uniprot ID: -

Gene ID: -

Swissprot:	P14151
Host Species:	Mouse
Isotype:	Mouse IgG1, κ
Isotype Control:	-
Conjugation:	-
Conjugation Information:	-
Buffer:	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.
Purification:	>98%, Protein A/G purified
Target:	CD62L
Cellular Localization:	-
Tissue Specificity:	-
Verified Samples:	Verified Samples in FCM: Human peripheral blood lymphocytes
Concentration:	≥ 1 mg/mL

Preparation & Storage

Storage: Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.

Shipping: Ice bag

Recommended Dilution: FCM 2 µg/mL(0.5×10⁶-1×10⁶ cells)

Recommended Usage:

Application	Recommended Usage
-	-

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes:

Centrifuge before opening to ensure complete recovery of vial contents.