

Purified Anti-Human CD25 Antibody [CHI621]

AGEL5094

Description

This Purified Anti-Human CD25 Antibody [CHI621] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL5094
Contents:	25µg, 100µg, 1mg Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	CHI621
Synonyms:	IMD, IDDM, IL2RA, CD25, IDDM10, IL2R, IMD41, TCGFR, p55, IL-2 receptor subunit alpha, IL-2R subunit alpha, IL2-RA, IL-2-RA, IL-2Rα, Interleukin-2 receptor subunit alpha, TAC antigen, Tac, p55, IL-2R&, alpha, alpha chain, CD25 antigen, IL 2 receptor alpha subunit, IL2 RA, Interleukin 2 receptor, Interleukin 2 receptor alpha, Interleukin 2 receptor alpha chain, t-cell growth factor receptor
Applications:	FCM
Reactivity:	Human
Immunogen:	Recombinant Human CD25 protein

Antibody Data

Uniprot ID:	-
Gene ID:	-
Swissprot:	P01589
Host Species:	Mouse

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Isotype:	Mouse IgG1, κ
Isotype Control:	-
Conjugation:	-
Conjugation Information:	-
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.
Purification:	>98%, Protein A/G purified
Target:	CD25
Cellular Localization:	-
Tissue Specificity:	-
Verified Samples:	Verified Samples in FCM: Human peripheral blood lymphocytes
Concentration:	≥ 1 mg/mL

Preparation & Storage

Storage: Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.

Shipping: Ice bag

Recommended Dilution: FCM 2 µg/mL(0.5×10⁶-1×10⁶ cells)

Recommended Usage:

Application	Recommended Usage
-	-

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.