

Purified Anti-Human CD192 Antibody [K036C2]

AGEL5129

Description

This Purified Anti-Human CD192 Antibody [K036C2] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL5129
Contents:	25µg, 100µg, 1mg Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	K036C2
Synonyms:	CKR, CMKBR, CC-CKR, CCR, CCR2, CC-CKR-2, CCR-2, CCR2A, CCR2B, CD192, CKR2, CKR2A, CKR2B, CMKBR2, MCP-1-R, FLJ78302, C C chemokine receptor type 2, C C CKR 2, CC chemokine receptor type 2, C-C chemokine receptor type 2, CC CKR 2, C-C CKR-2, CCCR2, CCR 2, CCR1L, CCR5L, CD192 antigen, Chemokine C C motif receptor 2, Chemokine CC Motif Receptor 2, CKR 2, cluster of differentiation 192, MCP 1 R, MCP1 RECEPTOR, MCP1R, MCP-1-R, Monocyte chemoattractant protein 1 receptor, Monocyte Chemotactic Protein 1 Receptor
Applications:	FCM
Reactivity:	Human
Immunogen:	Recombinant Human CD192 protein

Antibody Data

Uniprot ID:	-
Gene ID:	-

Swissprot:	P41597
Host Species:	Mouse
Isotype:	Mouse IgG2a, κ
Isotype Control:	-
Conjugation:	-
Conjugation Information:	-
Buffer:	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.
Purification:	>98%, Protein A/G purified
Target:	CD192
Cellular Localization:	-
Tissue Specificity:	-
Verified Samples:	Verified Samples in FCM: Human peripheral blood monocytes cell
Concentration:	≥ 1 mg/mL

Preparation & Storage

Storage: Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.

Shipping: Ice bag

Recommended Dilution: FCM 2 µg/mL(0.5×10⁶-1×10⁶ cells)

Recommended Usage:

Application	Recommended Usage
-	-

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.

