

Anti-Human CD14 Antibody [M5E2] -Low Endotoxin, Azide Free

AGEL5140

Description

This Anti-Human CD14 Antibody [M5E2] -Low Endotoxin, Azide Free is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	AGEL5140
Contents:	25µg, 100µg, 1mg Bradford Reagent: 1 vial (2ml)
Category:	Monoclonal Antibody
Clonality:	Monoclonal
Clone:	M5E2
Synonyms:	Monocyte Differentiation Antigen CD, CD14,Soluble, CD14 antigen, CD14 molecule, Monocyte Differentiation Antigen CD14, Myeloid cell-specific leucine-rich glycoprotein, LPS R, CD 14, CD_antigen=CD14, LPS-R, membrane-bound form, Mo2, Monocyte Differentiation Antigen CD14 urinary form, Myeloid cell specific leucine rich glycoprotein
Applications:	FCM
Reactivity:	Human
Immunogen:	Recombinant Human CD14 protein

Antibody Data

Uniprot ID:	-
Gene ID:	-
Swissprot:	P08571
Host Species:	Mouse

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Isotype:	Mouse IgG2a, κ
Isotype Control:	-
Conjugation:	-
Conjugation Information:	-
Buffer:	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.
Purification:	>98%, Protein A/G purified
Target:	CD14
Cellular Localization:	-
Tissue Specificity:	-
Verified Samples:	Verified Samples in FCM: Human peripheral blood monocytes
Concentration:	≥ 1 mg/mL

Preparation & Storage

Storage: Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.

Shipping: Ice bag

Recommended Dilution: FCM 2 µg/mL(0.5×10⁶-1×10⁶ cells)

Recommended Usage:

Application	Recommended Usage
-	-

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.