

# Purified Anti-Mouse CD8a Antibody [YTS 105.18]

AGEL5190

## Description

---

This Purified Anti-Mouse CD8a Antibody [YTS 105.18] is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	AGEL5190
<b>Contents:</b>	25µg, 100µg, 1mg Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Monoclonal Antibody
<b>Clonality:</b>	Monoclonal
<b>Clone:</b>	YTS 105.18
<b>Synonyms:</b>	T-cell surface glycoprotein Lyt, T-lymphocyte differentiation antigen T8/Leu, Lyt, Ly-2, Ly-B, Ly-35, Lyt-2, BB154331, CD antigen CD8a, MAL, T-cell surface glycoprotein CD8 alpha chain, T-cell surface glycoprotein Lyt-2, T-lymphocyte differentiation antigen T8/Leu-2, Lyt2, Cd8a, T8, Lyt-2, Ly-2, CD8
<b>Applications:</b>	<b>FCM</b>
<b>Reactivity:</b>	Mouse
<b>Immunogen:</b>	Recombinant Mouse CD8a protein

## Antibody Data

---

<b>Uniprot ID:</b>	-
<b>Gene ID:</b>	-
<b>Swissprot:</b>	P01731
<b>Host Species:</b>	Rat

<b>Isotype:</b>	Rat IgG2b, κ
<b>Isotype Control:</b>	-
<b>Conjugation:</b>	-
<b>Conjugation Information:</b>	-
<b>Buffer:</b>	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.
<b>Purification:</b>	>98%, Protein A/G purified
<b>Target:</b>	CD8a
<b>Cellular Localization:</b>	-
<b>Tissue Specificity:</b>	-
<b>Verified Samples:</b>	Verified Samples in FCM: C57/BL6 Mouse splenocytes
<b>Concentration:</b>	≥ 1 mg/mL

## Preparation & Storage

---

**Storage:** Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.

**Shipping:** Ice bag

**Recommended Dilution:** FCM 2 µg/mL(0.5×10<sup>6</sup>-1×10<sup>6</sup> cells)

**Recommended Usage:**

Application	Recommended Usage
-	-

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.