

# Anti-WTAP [R09-9D8] Monoclonal Antibody

## AGMB00750

### Description

---

This Anti-WTAP [R09-9D8] Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

<b>SKU:</b>	AGMB00750
<b>Contents:</b>	20µl, 50µl, 100µl Bradford Reagent: 1 vial (2ml)
<b>Synonyms:</b>	Mum2, WTAP, hFL(2)D, WT1-associated protein, Wilms tumor 1-associating protein, Pre-mRNA-splicing regulator WTAP
<b>Applications:</b>	<b>WB</b> <b>IHC-P</b> <b>ICC/IF</b> <b>FC</b> <b>IP</b>
<b>Research Area:</b>	Epigenetics
<b>Form:</b>	Liquid

### Antibody Data

---

<b>Reactivity:</b>	Human
<b>Clone:</b>	R09-9D8
<b>Clonality:</b>	Monoclonal Antibody
<b>SwissProt ID:</b>	Q15007
<b>Immunogen:</b>	Recombinant protein of human WTAP
<b>Gene ID:</b>	9589

#### Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

**Gene Name:** WTAP  
**Host Species:** Rabbit  
**Isotype:** IgG  
**Purification:** Affinity Purified  
**Conjugated:** Unconjugated  
**Modification:** Unmodified  
**Molecular Weight:** Calculated MW: 44 kDa, Observed MW: 55 kDa

## Preparation & Storage

---

**Storage:** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.  
 Store Bradford Reagent at Room Temperature for 1 Year.

**Storage Buffer:** Liquid in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA.

<b>Antibody Dilution Ratio:</b>	<b>Application</b>	<b>Antibody Dilution Ratio</b>
	WB	1:500-1:1000
	IHC	1:50-1:100
	ICC/IF	1:200-1:500
	FC	1:200-1:500
	IP	1:20-1:50

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol.