

## Map2 Antibody

CAB0453

### Description

---

This Map2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB0453  
**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** MAP-2, MAP2A, MAP2B, MAP2C, MAP2  
**Clone:** -  
**Applications:** **WB** | **IHC-P** | **ELISA** | **IF-P**  
**Conjugation:** Unconjugated  
**Reactivity:** Mouse, Rat

### Antibody Data

---

**Gene ID:** 4133  
**Uniprot:** AB\_2757199  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 220-280kd(MAP2A, MAP2B)  
**Calculated MW:** 200kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.


**Positive Sample:** Mouse brain

<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:100 - 1:500</td></tr> <tr> <td><b>IF-P</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>IHC-P</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:100 - 1:500	<b>IF-P</b>	1:50 - 1:200	<b>IHC-P</b>	1:50 - 1:200	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:100 - 1:500								
<b>IF-P</b>	1:50 - 1:200								
<b>IHC-P</b>	1:50 - 1:200								
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data


---




Western blot analysis of various lysates, using Rabbit pAb (CAB0453) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Rabbit pAb (CAB0453) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of paraffin-embedded mouse brain using Rabbit pAb (CAB0453) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded rat brain using Rabbit pAb (CAB0453) at dilution of 1:20 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.