

UCK2 Antibody

CAB10815

Description

This UCK2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB10815
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: UK, UMPK, TSA903, UCK2
Clone: -
Applications: WB ELISA
Conjugation: Unconjugated
Reactivity: Human, Mouse


Antibody Data

Gene ID: 7371
Uniprot: AB_2758243
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 29kDa
Calculated MW: 29kDa

Preparation & Storage

| | | | | | |
|---|--|-----------|----------------|--------------|---|
| Storage: | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. | | | | |
| | Store Bradford Reagent at Room Temperature for 1 Year. | | | | |
| Positive Sample: | A-549, HeLa, 293T, B-cell, Mouse thymus | | | | |
| Recommended Dilutions: | <table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | WB | 1:500 - 1:2000 | ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| WB | 1:500 - 1:2000 | | | | |
| ELISA | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. | | | | |
| Protein Quantification (Optional): | To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol | | | | |

Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB10815) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 90s.