

F11R Antibody

CAB1241

Description

This F11R Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:

CAB1241

Contents:

20 μ L, 100 μ L

Bradford Reagent: 1 vial (2ml)

Category:

Polyclonal Antibody

Synonyms:

JAM, KAT, JAM1, JAMA, JCAM, CD321, PAM-1, JAM-A/CD321/F11R

Clone:

-

Applications:

WB **IHC-P** **ELISA**

Conjugation:

Unconjugated

Reactivity:

Human, Mouse, Rat

Antibody Data

Gene ID:

50848

Uniprot:

AB_2759253

Host Species:

Rabbit

Purification:

Affinity purification

Observed MW:

37kDa

Calculated MW:

33kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: HepG2, SKOV3, Mouse lung, Rat lung


Recommended Dilutions:	WB	1:500 - 1:1000
	IHC-P	1:50 - 1:200
	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Western blot analysis of various lysates using JAM-A/F11R Rabbit pAb (CAB1241) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded Human esophageal cancer using JAM-A//F11R Rabbit pAb (CAB1241) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta using JAM-A//F11R Rabbit pAb (CAB1241) at dilution of 1:50 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.