

## WNT5A Antibody

CAB12744

### Description

---

This WNT5A Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB12744  
**Contents:** 20  $\mu$ L, 100  $\mu$ L  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** hWNT5A, WNT5A  
**Clone:** -  
**Applications:** **WB** | **IHC-P** | **IF/ICC** | **ELISA**  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Mouse, Rat

### Antibody Data

---

**Gene ID:** 7474  
**Uniprot:** AB\_2759588  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 45kDa  
**Calculated MW:** 42kDa

## Preparation & Storage

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** SKOV3, Rat lung


|                               |   |           |                |              |               |               |              |              |   |
|-------------------------------|---|-----------|----------------|--------------|---------------|---------------|--------------|--------------|---|
| <b>Recommended Dilutions:</b> | <table border="1"> <tr> <td><b>WB</b></td><td>1:500 - 1:1000</td></tr> <tr> <td><b>IHC-P</b></td><td>1:100 - 1:500</td></tr> <tr> <td><b>IF/ICC</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table> | <b>WB</b> | 1:500 - 1:1000 | <b>IHC-P</b> | 1:100 - 1:500 | <b>IF/ICC</b> | 1:50 - 1:200 | <b>ELISA</b> | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |
| <b>WB</b>                     | 1:500 - 1:1000  |           |                |              |               |               |              |              |   |
| <b>IHC-P</b>                  | 1:100 - 1:500   |           |                |              |               |               |              |              |   |
| <b>IF/ICC</b>                 | 1:50 - 1:200  |           |                |              |               |               |              |              |   |
| <b>ELISA</b>                  | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.   |           |                |              |               |               |              |              |   |

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


## Validation Data




Western blot analysis of various lysates using (CAB12744) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Human thyroid cancer tissue using WNT5A Rabbit pAb (CAB12744) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human placenta tissue using WNT5A Rabbit pAb (CAB12744) at a dilution of 1:300 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of PC-12 cells using WNT5A Rabbit pAb (CAB12744) at dilution of 1:50 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.