

IFNL3 Antibody

CAB12908

Description

This IFNL3 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB12908

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: IL28B, IL28C, IL-28B, IL-28C, IFN-lambda-3, IFN-lambda-4, IFNL3

Clone: -

Applications: IHC-P ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 282617

Uniprot: AB_2759753

Host Species: Rabbit

Purification: Affinity purification

Observed MW: -

Calculated MW: 22kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample:


-

Recommended Dilutions:


IHC-P	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Immunohistochemistry analysis of paraffin-embedded Rat brain using IFNL3 Rabbit pAb (CAB12908) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human stomach using IFNL3 Rabbit pAb (CAB12908) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse lung using IFNL3 Rabbit pAb (CAB12908) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.