

DKC1 Antibody

CAB12914

Description

This DKK1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB12914
Contents: 20 μ L, 100 μ L
Bradford Reagent: 1 vial (2ml)
Category: Polyclonal Antibody
Synonyms: DKK, CBF5, DKKX, NAP57, NOLA4, XAP101, DKK1
Clone: -
Applications: **WB** **IHC-P** **IF/ICC** **ELISA**
Conjugation: Unconjugated
Reactivity: Human, Mouse


Antibody Data

Gene ID: 1736
Uniprot: AB_2759759
Host Species: Rabbit
Purification: Affinity purification
Observed MW: 57kDa
Calculated MW: 58kDa


Preparation & Storage

Storage:	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.								
	Store Bradford Reagent at Room Temperature for 1 Year.								
Positive Sample:	HeLa, 22Rv1, HepG2, MCF7, K-562								
Recommended Dilutions:	<table border="1"> <tr> <td>WB</td><td>1:500 - 1:2000</td></tr> <tr> <td>IHC-P</td><td>1:50 - 1:100</td></tr> <tr> <td>IF/ICC</td><td>1:50 - 1:100</td></tr> <tr> <td>ELISA</td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	WB	1:500 - 1:2000	IHC-P	1:50 - 1:100	IF/ICC	1:50 - 1:100	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
WB	1:500 - 1:2000								
IHC-P	1:50 - 1:100								
IF/ICC	1:50 - 1:100								
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol								


Validation Data



Western blot analysis of various lysates using Rabbit pAb (CAB12914) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Rabbit pAb (CAB12914) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of U-2 OS cells using Rabbit pAb (CAB12914) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.