

## Septin 5 Antibody

**CAB12953**

### Description

---

This Septin 5 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

<b>SKU:</b>	CAB12953
<b>Contents:</b>	20 µL, 100 µL Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	Polyclonal Antibody
<b>Synonyms:</b>	H5, SEPT5, CDCREL, PNUTL1, CDCREL1, CDCREL-1, HCDCREL-1, Septin 5
<b>Clone:</b>	-
<b>Applications:</b>	<b>WB</b> <b>IHC-P</b> <b>IF/ICC</b> <b>ELISA</b>
<b>Conjugation:</b>	Unconjugated
<b>Reactivity:</b>	Human, Mouse, Rat

### Antibody Data

---

<b>Gene ID:</b>	5413
<b>Uniprot:</b>	AB_2759799
<b>Host Species:</b>	Rabbit
<b>Purification:</b>	Affinity purification
<b>Observed MW:</b>	43kDa
<b>Calculated MW:</b>	43kDa


## Preparation & Storage

---


<b>Storage:</b>	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3. Store Bradford Reagent at Room Temperature for 1 Year.								
<b>Positive Sample:</b>	Mouse brain, Rat brain								
<b>Recommended Dilutions:</b>	<table border="1"> <tr> <td><b>WB</b></td><td>1:1000 - 1:5000</td></tr> <tr> <td><b>IHC-P</b></td><td>1:100 - 1:200</td></tr> <tr> <td><b>IF/ICC</b></td><td>1:50 - 1:200</td></tr> <tr> <td><b>ELISA</b></td><td>Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.</td></tr> </table>	<b>WB</b>	1:1000 - 1:5000	<b>IHC-P</b>	1:100 - 1:200	<b>IF/ICC</b>	1:50 - 1:200	<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
<b>WB</b>	1:1000 - 1:5000								
<b>IHC-P</b>	1:100 - 1:200								
<b>IF/ICC</b>	1:50 - 1:200								
<b>ELISA</b>	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.								
<b>Protein Quantification (Optional):</b>	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <a href="https://www.assaygenie.com/bradford-protein-assay-protocol/">https://www.assaygenie.com/bradford-protein-assay-protocol/</a> to view the full protocol								

## Validation Data


---



Western blot analysis of various lysates, using Septin 5 Rabbit pAb (CAB12953) at 1:3000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Septin 5 Rabbit pAb (CAB12953) at dilution of 1:100 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.



Immunofluorescence analysis of U-2 OS cells using Septin 5 Rabbit pAb (CAB12953) at dilution of 1:100. Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.