

## UPF2 Antibody

**CAB13411**

### Description

---

This UPF2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

**SKU:** CAB13411  
**Contents:** 20 µL, 100 µL  
Bradford Reagent: 1 vial (2ml)  
**Category:** Polyclonal Antibody  
**Synonyms:** HUPF2, RENT2, smg-3, UPF2  
**Clone:** -  
**Applications:** **WB** | **IHC-P** | **IP** | **ELISA**  
**Conjugation:** Unconjugated  
**Reactivity:** Human, Rat

### Antibody Data

---

**Gene ID:** 26019  
**Uniprot:** AB\_2760273  
**Host Species:** Rabbit  
**Purification:** Affinity purification  
**Observed MW:** 170kDa  
**Calculated MW:** 148kDa

## Preparation & Storage

---

**Storage:** Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

Store Bradford Reagent at Room Temperature for 1 Year.

**Positive Sample:** Jurkat, 293T


**Recommended Dilutions:**

|              |   |
|--------------|---|
| <b>WB</b>    | 1:500 - 1:2000  |
| <b>IHC-P</b> | 1:50 - 1:200  |
| <b>IP</b>    | 0.5µg-4µg antibody for 200µg-400µg extracts of whole cells  |
| <b>ELISA</b> | Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements. |


**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---



Western blot analysis of various lysates, using Rabbit pAb (CAB13411) at 1:2000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded Rat kidney using Rabbit pAb (CAB13411) at dilution of 1:200 (40x lens). Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to IHC staining.