

CLU Antibody

CAB13479

Description

This CLU Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: CAB13479

Contents: 20 µL, 100 µL

Bradford Reagent: 1 vial (2ml)

Category: Polyclonal Antibody

Synonyms: CLI, AAG4, APOJ, CLU1, CLU2, KUB1, SGP2, APO-J, SGP-2, SP-40, TRPM2, TRPM-2, NA1/NA2, Clusterin

Clone: -

Applications: WB IHC-P IF/ICC ELISA

Conjugation: Unconjugated

Reactivity: Human, Mouse, Rat

Antibody Data

Gene ID: 1191

Uniprot: AB_2760341

Host Species: Rabbit

Purification: Affinity purification

Observed MW: 38kDa/57kDa

Calculated MW: 52kDa

Preparation & Storage

Storage: Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.


Store Bradford Reagent at Room Temperature for 1 Year.

Positive Sample: HepG2, U-251MG, Human plasma, Mouse plasma


Recommended Dilutions:	WB	1:500 - 1:2000
	IHC-P	1:50 - 1:200
	IF/ICC	1:50 - 1:200
	ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data



Western blot analysis of various lysates using Clusterin Rabbit pAb (CAB13479) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (CABS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (AbGn00020). Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Human colon cancer tissue using Clusterin Rabbit pAb (CAB13479) at a dilution of 1:100 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate Buffer(pH 6.0) prior to IHC staining.



Immunofluorescence analysis of U2OS cells using Clusterin Rabbit pAb (CAB13479). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (CABS007) at 1:500 dilution. Blue: DAPI for nuclear staining.